

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/12, 5/10, C07K 14/47, 14/705, 16/18, 16/28, C12Q 1/68</b>		<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/01821</b> <b>(43) International Publication Date:</b> 13 January 2000 (13.01.00)
<b>(21) International Application Number:</b> PCT/US99/15121 <b>(22) International Filing Date:</b> 2 July 1999 (02.07.99) <b>(30) Priority Data:</b> 60/091,677 2 July 1998 (02.07.98) US <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 60/091,677 (CIP) Filed on 2 July 1998 (02.07.98) <b>(71) Applicant (for all designated States except US):</b> INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue #30, Mountain View, CA 94040 (US). GUEGLER, Karl, J. [CH/US]; 1048 Oakland Avenue, Menlo Park, CA 94025 (US). GORGONE, Gina, A. [US/US]; 1253 Pinecrest			<b>(74) Agents:</b> BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> NEUROTRANSMISSION ASSOCIATED PROTEINS			
<b>(57) Abstract</b> <p>The invention provides human neurotransmission associated proteins (NTAP) and polynucleotides which identify and encode NTAP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing disorders associated with expression of NTAP.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## NEUROTRANSMISSION ASSOCIATED PROTEINS

### TECHNICAL FIELD

5           This invention relates to nucleic acid and amino acid sequences of neurotransmission associated proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cancer and immune and neurological disorders.

### BACKGROUND OF THE INVENTION

10           Neurotransmission is the basic process of transmitting nerve signals, in the form of electrical and chemical impulses, between nerve cells which carry signals from sensory receptors (e.g., for light, touch, pressure, odor and taste) to the brain, and similar signals from the brain to effector targets or organs such as skeletal or smooth muscle. The nervous system is composed of the central nervous system (CNS), consisting of the brain and spinal cord, and the peripheral  
15   nervous system (PNS), consisting of afferent neural pathways for conducting nerve impulses from sensory organs to the CNS, and efferent neural pathways for conducting motor impulses from the CNS to effector organs. The PNS can be further divided into the somatic nervous system, which regulates voluntary motor activity such as for skeletal muscle, and the autonomic nervous system, which regulates involuntary motor activity for internal organs such as the heart, lungs, and viscera.

20           The basic cellular unit of neurotransmission is the nerve cell, or neuron. The process of neurotransmission involves the transmission of a neural signal between neurons by means of a combination of electrical and chemical impulses. The process begins with the stimulation of a nerve and the generation of an electrical impulse at which travels along the axon of the neuron to its terminus. Neurons are separated from one another by a space (the synapse or synaptic cleft)  
25   which must be bridged to transmit the signal to another neuron. This accomplished by a specialized form of vesicle transport which uses a neurotransmitter signaling molecule stored in a membrane-bound synaptic vesicle at the terminus of the neuron. A change in electrical potential at the nerve terminal resulting from the electrical impulse triggers the release of the neurotransmitter from the synaptic vesicle by exocytosis. The neurotransmitter rapidly diffuses  
30   across the synaptic cleft separating the presynaptic nerve cell from the postsynaptic cell and provokes a change in electrical potential in the latter by binding to receptors and opening transmitter-gated ion channels located in the plasma membrane of the postsynaptic cell. The change in membrane potential of the postsynaptic cell may serve either to excite or inhibit further transmission of the nerve impulse.

Neurotransmitters comprise a diverse group of some 30 substances which include acetylcholine, monoamines such as serotonin, dopamine, and histamine, and amino acids such as gamma-aminobutyric acid (GABA), glutamate, and aspartate, and neuropeptides such as endorphins and enkephalins. (McCance, K.L. and Huether, S.E. (1994) PATHOPHYSIOLOGY,  
5 The Biologic Basis for Disease in Adults and Children, 2nd edition, Mosby, St. Louis, MO, pp 403-404.) Many of these molecules have more than one function and the effects may be excitatory, e.g. to depolarize the postsynaptic cell plasma membrane and stimulate nerve impulse transmission, or inhibitory, e.g. to hyperpolarize the plasma membrane and inhibit nerve impulse transmission.

10 Neurotransmitters and their receptors are targets of pharmacological agents aimed at controlling neurological function. For example GABA is the major inhibitory neurotransmitter in the CNS, and GABA receptors are the principal target of sedatives such as benzodiazepines and barbiturates which act by enhancing GABA mediated effects. (Katzung, B.G. (1995) Basic and Clinical Pharmacology, 6th edition, Appleton & Lange, Norwalk, CT, pp. 338-339) Aberrant  
15 activity of neurotransmitters and their receptors are involved in various neurological conditions. Alzheimer's disease is associated with a decrease in acetylcholine-secreting neurons, and myasthenia gravis results from a reduction in acetylcholine receptors. Destruction of dopamine-secreting receptors and overstimulation of N-methyl-D-aspartate (NMDA) receptors in the brain is implicated in neuronal cell death associated with disorders such as stroke, epilepsy, Parkinson's  
20 disease and Alzheimer's disease. (Planells-Cases, R. Et al. (1993) Proc. Natl. Acad. Sci. USA 90:5057-5061.)

Various molecules are associated with synaptic vesicles that store and transport neurotransmitters. Synaptic vesicles of mature neurons have been shown to possess a specific complement of membrane proteins which are restricted to these vesicles, and at least 15 synaptic  
25 vesicle proteins have been characterized. (Sudhof, T.C. and Jahn, R. (1991) Neuron 6:665-677.) Synaptophysin and synaptogyrin are two such proteins that colocalize in synaptic vesicle preparations. (Stenius, K. et al. (1995) J. Cell Biology 131:1801-1809.) The precise functions of these proteins in neurosecretion are unknown. Both have four transmembrane domains and are highly expressed in neuronal tissues, but may have nonneuronal isoforms as well. Syntaxins are  
30 another family of proteins which are involved in synaptic vesicle transport, are associated with the plasma membrane of the neuron, and function as recognition sites for docking of the synaptic vesicle with the plasma membrane. Syntaxin interacts with complementary proteins on the synaptic vesicle (v-SNARES) to form a complex that initiates fusion of the vesicle with the plasma membrane prior to release of the neurotransmitter into the synapse. (Bock, J.B. et al. (1996) J.  
35 Biol. Chem. 271:17961-17965.)

Various proteins are also associated with the sensory response to stimuli in organs that trigger a nerve signal. Rhodopsin is the photosensitive protein in rod cells of the eye that undergoes a conformational change during the absorption of light by an associated chromophore, retinal. This conformational change initiates a photochemical cascade that leads to a nerve signal.

- 5 Odorant detection is mediated by receptor neurons in the olfactory mucosa and also involves distinct odorant-binding proteins that act to bind and carry specific odorant molecules across the mucus layer to odorant receptor sites. (Dear, T.N. et al. (1991) EMBO Journal 10:2813-2819.)

The discovery of new neurotransmission associated proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the  
10 diagnosis, treatment, and prevention of cancer and immune and neurological disorders.

## SUMMARY OF THE INVENTION

The invention features substantially purified polypeptides, neurotransmission associated proteins referred to collectively as "NTAP". In one aspect, the invention provides a substantially  
15 purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6 (SEQ ID NO:1-6), and fragments thereof.

The invention further provides a substantially purified variant having at least 90% amino acid identity to the amino acid sequences of SEQ ID NO:1-6, or to a fragment of any of these  
20 sequences. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting  
25 of SEQ ID NO:1-6, and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof, as well as an isolated and purified polynucleotide having a sequence which is  
30 complementary to the polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12 (SEQ ID NO:7-12), and fragments

thereof. The invention further provides an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide sequence comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:7-12, and fragments thereof, as well as an isolated and purified polynucleotide having a sequence which is

- 5 complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:7-12, and fragments thereof.

The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof. In another aspect, the expression  
10 vector is contained within a host cell.

The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing at least a fragment of a polynucleotide encoding the polypeptide under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

- 15 The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and  
20 fragments thereof, as well as a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a cancer, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of the polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof.

- 25 The invention also provides a method for treating or preventing an immune disorder, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of the polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof.

The invention also provides a method for treating or preventing a neurological disorder,  
30 the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-6, and fragments thereof.

The invention also provides a method for detecting a polynucleotide, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence encoding  
35 the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID

NO:1-6, and fragments thereof to at least one of the nucleic acids of the biological sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the biological sample. In one aspect, the method further comprises amplifying the polynucleotide  
 5 prior to hybridization.

### BRIEF DESCRIPTION OF THE TABLES

In Table 1, columns 1 and 2 show the sequence identification numbers (SEQ ID NO:) of the amino acid and nucleic acid sequence, respectively. Column 3 shows the Clone ID of the  
 10 Incyte Clone in which nucleic acids encoding each NTAP were first identified, and column 4, the cDNA library of this clone. Column 5 is entitled fragments, and shows the Incyte clones (and libraries) and shotgun sequences useful as fragments in hybridization technologies, and which are part of the consensus nucleotide sequence of each NTAP.

The columns of table 2 show various properties of the polypeptides of the invention:  
 15 column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues; column 3, potential phosphorylation sites; column 4, potential glycosylation sites; column 5, signature sequences associated with known proteins; column 6, the identity of the protein; and column 7, analytical methods used to identify the protein through sequence homologies and protein motifs.

The columns of table 3 show the tissue expression of each nucleic acid sequence by  
 20 northern analysis, diseases or disorders associated with this tissue expression, and the vector into which each cDNA was cloned.

Table 4 shows the SEQ ID NO:, Incyte clone number and the associated library in which nucleic acid sequences encoding NTAP were first identified, and a brief description of the library.

Table 5 shows the programs/algorithms, descriptions, references and threshold parameters  
 25 used to identify and characterize NTAP.

### DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described, as these may vary. It is also to be understood that the terminology  
 30 used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an

antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

## DEFINITIONS

"NTAP," as used herein, refers to the amino acid sequences, or variant thereof, of substantially purified NTAP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and preferably the human species, from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist," as used herein, refers to a molecule which, when bound to NTAP, increases or prolongs the duration of the effect of NTAP. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of NTAP.

An "allelic variant," as this term is used herein, is an alternative form of the gene encoding NTAP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding NTAP, as described herein, include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide the same as NTAP or a polypeptide with at least one functional characteristic of NTAP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding NTAP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding NTAP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues

which produce a silent change and result in a functionally equivalent NTAP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of NTAP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and phenylalanine and tyrosine.

The terms "amino acid" or "amino acid sequence," as used herein, refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. In this context, "fragments," "immunogenic fragments," or "antigenic fragments" refer to fragments of NTAP which are preferably at least 5 to about 15 amino acids in length, most preferably at least 14 amino acids, and which retain some biological activity or immunological activity of NTAP. Where "amino acid sequence" is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification," as used herein, relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art. (See, e.g., Dieffenbach, C.W. and G.S. Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, pp.1-5.)

The term "antagonist," as it is used herein, refers to a molecule which, when bound to NTAP, decreases the amount or the duration of the effect of the biological or immunological activity of NTAP. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies, or any other molecules which decrease the effect of NTAP.

As used herein, the term "antibody" refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies that bind NTAP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant," as used herein, refers to that fragment of a molecule

(i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (given regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact  
5 antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense," as used herein, refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced  
10 by the cell to form duplexes and to block either transcription or translation. The designation "negative" can refer to the antisense strand, and the designation "positive" can refer to the sense strand.

As used herein, the term "biologically active," refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise,  
15 "immunologically active" refers to the capability of the natural, recombinant, or synthetic NTAP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" or "complementarity," as used herein, refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" binds to the  
20 complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in  
25 amplification reactions, which depend upon binding between nucleic acids strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" or a "composition comprising a given amino acid sequence," as these terms are used herein, refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may  
30 comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding NTAP or fragments of NTAP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts, e.g., NaCl, detergents, e.g., sodium dodecyl sulfate (SDS), and other components, e.g.,  
35 Denhardt's solution, dry milk, salmon sperm DNA, etc.

“Consensus sequence,” as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using XL-PCR™ kit (Perkin-Elmer, Norwalk, CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte Clone using a computer program for fragment  
5 assembly, such as the GELVIEW™ Fragment Assembly system (GCG, Madison, WI). Some sequences have been both extended and assembled to produce the consensus sequence.

As used herein, the term “correlates with expression of a polynucleotide” indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding NTAP, by Northern analysis is indicative of the presence of nucleic acids encoding NTAP in a  
10 sample, and thereby correlates with expression of the transcript from the polynucleotide encoding NTAP.

A “deletion,” as the term is used herein, refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term “derivative,” as used herein, refers to the chemical modification of a polypeptide  
15 sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or  
20 immunological function of the polypeptide from which it was derived.

The term “similarity,” as used herein, refers to a degree of complementarity. There may be partial similarity or complete similarity. The word “identity” may substitute for the word “similarity.” A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as “substantially similar.” The  
25 inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of  
30 reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence  
35 or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" or "% identity" refer to the percentage of sequence similarity found in a comparison of two or more amino acid or nucleic acid sequences. Percent identity can be determined electronically, e.g., by using the MegAlign™ program (DNASTAR, Inc., Madison WI). The MegAlign™ program can create alignments between two or more sequences according to different methods, e.g., the clustal method. (See, e.g., Higgins, D.G. and P.M. Sharp (1988) *Gene* 73:237-244.) The clustal algorithm groups sequences into clusters by examining the distances between all pairs. The clusters are aligned pairwise and then in groups. The percentage similarity between two amino acid sequences, e.g., sequence A and sequence B, is calculated by dividing the length of sequence A, minus the number of gap residues in sequence A, minus the number of gap residues in sequence B, into the sum of the residue matches between sequence A and sequence B, times one hundred. Gaps of low or of no similarity between the two amino acid sequences are not included in determining percentage similarity. Percent identity between nucleic acid sequences can also be counted or calculated by other methods known in the art, e.g., the Jotun Hein method. (See, e.g., Hein, J. (1990) *Methods Enzymol.* 183:626-645.) Identity between sequences can also be determined by other methods known in the art, e.g., by varying hybridization conditions.

"Human artificial chromosomes" (HACs), as described herein, are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance. (See, e.g., Harrington, J.J. et al. (1997) *Nat Genet.* 15:345-355.)

The term "humanized antibody," as used herein, refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization," as the term is used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

As used herein, the term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_0t$  or  $R_0t$  analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" or "addition," as used herein, refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, to the sequence found in the naturally occurring molecule.

"Immune response" can refer to conditions associated with inflammation, trauma, immune

disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term "microarray," as used herein, refers to an arrangement of distinct  
5 polynucleotides arrayed on a substrate, e.g., paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support.

The terms "element" or "array element" as used herein in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate," as it appears herein, refers to a change in the activity of NTAP. For  
10 example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of NTAP.

The phrases "nucleic acid" or "nucleic acid sequence," as used herein, refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded  
15 and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material. In this context, "fragments" refers to those nucleic acid sequences which, comprise a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:7-12, for example, as distinct from any other sequence in the same genome. For example, a fragment of SEQ ID NO:7-12 is useful in hybridization and amplification technologies  
20 and in analogous methods that distinguish SEQ ID NO:7-12 from related polynucleotide sequences. A fragment of SEQ ID NO:7-12 is at least about 15-20 nucleotides in length. The precise length of the fragment of SEQ ID NO:7-12 and the region of SEQ ID NO:7-12 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment. In some cases, a fragment, when translated, would produce  
25 polypeptides retaining some functional characteristic, e.g., antigenicity, or structural domain characteristic, e.g., ATP-binding site, of the full-length polypeptide.

The terms "operably associated" or "operably linked," as used herein, refer to functionally related nucleic acid sequences. A promoter is operably associated or operably linked with a coding sequence if the promoter controls the translation of the encoded polypeptide. While  
30 operably associated or operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements, e.g., repressor genes, are not contiguously linked to the sequence encoding the polypeptide but still bind to operator sequences that control expression of the polypeptide.

The term "oligonucleotide," as used herein, refers to a nucleic acid sequence of at least  
35 about 6 nucleotides to 60 nucleotides, preferably about 15 to 30 nucleotides, and most preferably

about 20 to 25 nucleotides, which can be used in PCR amplification or in a hybridization assay or microarray. As used herein, the term "oligonucleotide" is substantially equivalent to the terms "amplimer," "primer," "oligomer," and "probe," as these terms are commonly defined in the art.

"Peptide nucleic acid" (PNA), as used herein, refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell. (See, e.g., Nielsen, P.E. et al. (1993) *Anticancer Drug Des.* 8:53-63.)

The term "sample," as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acids encoding NTAP, or fragments thereof, or NTAP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a solid support; a tissue; a tissue print; etc.

As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

As used herein, the term "stringent conditions" refers to conditions which permit hybridization between polynucleotides and the claimed polynucleotides. Stringent conditions can be defined by salt concentration, the concentration of organic solvent, e.g., formamide, temperature, and other conditions well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

The term "substantially purified," as used herein, refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

A "substitution," as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Transformation," as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions

according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle

- 5 bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

- A "variant" of NTAP polypeptides, as used herein, refers to an amino acid sequence that  
10 is altered by one or more amino acid residues. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have "nonconservative" changes (e.g., replacement of glycine with tryptophan). Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be  
15 substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, LASERGENE™ software (DNASTAR).

- The term "variant," when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to NTAP. This definition may also include, for  
20 example, "allelic" (as defined above), "splice," "species," or "polymorphic" variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains. Species variants are polynucleotide sequences that vary from one species to another. The resulting  
25 polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for  
30 a disease state.

## THE INVENTION

The invention is based on the discovery of new human neurotransmission associated proteins (NTAP), the polynucleotides encoding NTAP, and the use of these compositions for the

diagnosis, treatment, or prevention of cancer and immune and neurological disorders.

Table 1 shows the Clone ID of the Incyte Clone in which nucleic acids encoding each NTAP were first identified, and the Incyte clones (and libraries) and shotgun sequences which are part of the consensus nucleotide sequence of each NTAP. Table 2 shows various properties of the polypeptides of the invention and methods used to identify the protein through sequence homologies and protein motifs. Table 3 shows the tissue expression of each nucleic acid sequence by northern analysis and diseases or disorders associated with this tissue expression.

The following represent unique fragments of the nucleotide sequences encoding NTAP and useful, for example, as hybridization probes: the fragment of SEQ ID NO:7 from about nucleotide 568 to about nucleotide 711; the fragment of SEQ ID NO:8 from about nucleotide 435 to about nucleotide 479; the fragment of SEQ ID NO:9 from about nucleotide 335 to about nucleotide 388; the fragment of SEQ ID NO:10 from about nucleotide 393 to about nucleotide 452; the fragment of SEQ ID NO:11 from about nucleotide 758 to about nucleotide 799; and the fragment of SEQ ID NO:12 from about nucleotide 325 to about nucleotide 378.

The invention also encompasses NTAP variants. A preferred NTAP variant is one which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% amino acid sequence identity to the NTAP amino acid sequence, and which contains at least one functional or structural characteristic of NTAP.

The invention also encompasses polynucleotides which encode NTAP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:7-12, which encodes an NTAP

The invention also encompasses a variant of a polynucleotide sequence encoding NTAP. In particular, such a variant polynucleotide sequence will have at least about 80%, more preferably at least about 90%, and most preferably at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding NTAP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:7-12 which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:7-12. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of NTAP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding NTAP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide

sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring NTAP, and all such variations are to be considered as being specifically disclosed.

5           Although nucleotide sequences which encode NTAP and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring NTAP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding NTAP possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of  
10 the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding NTAP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

15           The invention also encompasses production of DNA sequences which encode NTAP and NTAP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding NTAP or any fragment thereof.

20           Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:7-12 or fragments thereof, under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) For example, stringent salt concentration will ordinarily be less than about 750 mM  
25 NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and most preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide, and most preferably at least about 50% formamide. Stringent temperature conditions will  
30 ordinarily include temperatures of at least about 30°C, more preferably of at least about 37°C, and most preferably of at least about 42°C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are accomplished by combining these various conditions as needed. In a preferred embodiment,  
35 hybridization will occur at 30°C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a

more preferred embodiment, hybridization will occur at 37°C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 µg/ml denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will occur at 42°C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50 % formamide, and 200 µg/ml ssDNA. Useful variations on these conditions

5 will be readily apparent to those skilled in the art.

The washing steps which follow hybridization can also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM

10 NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include temperature of at least about 25°C, more preferably of at least about 42°C, and most preferably of at least about 68°C. In a preferred embodiment, wash steps will occur at 25°C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at

15 42°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a most preferred embodiment, wash steps will occur at 68°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. Additional variations on these conditions will be readily apparent to those skilled in the art.

Methods for DNA sequencing and analysis are well known in the art. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE® (Amersham

20 Pharmacia Biotech Ltd., Uppsala, Sweden), Taq polymerase (Perkin-Elmer ), thermostable T7 polymerase (Amersham Pharmacia Biotech Ltd., Uppsala, Sweden), or combinations of polymerases and proofreading exonucleases, such as those found in the ELONGASE™ amplification system (Life Technologies, Inc., Rockville, MD). Preferably, sequence preparation is automated with machines, e.g., the ABI CATALYST™ 800 (Perkin-Elmer ) or MICROLAB®

25 2200 (Hamilton Co., Reno, NV) systems, in combination with thermal cyclers. Sequencing can also be automated, such as by ABI PRISM™ 373 or 377 systems (Perkin-Elmer ) or the MEGABACE™ 1000 capillary electrophoresis system (Molecular Dynamics, Inc., Sunnyvale, CA). Sequences can be analyzed using computer programs and algorithms well known in the art. (See, e.g., Ausubel, F. M. et al. (1997) Current Protocols in Molecular Biology, John Wiley &

30 Sons, New York, NY, unit 7.7; and Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, Inc, New York, NY.)

The nucleic acid sequences encoding NTAP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be

35 employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence

from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR.

Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-306). Additionally, one may use PCR, nested primers, and PromoterFinder™ libraries (Clontech, Palo Alto, CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO™ 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., Genotyper™ and Sequence Navigator™ (Perkin-Elmer Corp.), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode NTAP may be cloned in recombinant DNA molecules that direct expression of NTAP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the

same or a functionally equivalent amino acid sequence may be produced and used to express NTAP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter NTAP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding NTAP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucl. Acids Res. Symp. Ser. 215-223, and Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232.) Alternatively, NTAP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A Peptide Synthesizer (Perkin-Elmer Corp.). Additionally, the amino acid sequence of NTAP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman and Co., New York, NY.)

In order to express a biologically active NTAP, the nucleotide sequences encoding NTAP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding NTAP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding NTAP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding NTAP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where

only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding NTAP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, ch. 4, 8, and 16-17; and Ausubel, F.M. et al. (1995, and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding NTAP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV)) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding NTAP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding NTAP can be achieved using a multifunctional E. coli vector such as Bluescript® (Stratagene) or pSport1™ plasmid (GIBCO BRL). Ligation of sequences encoding NTAP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When large quantities of NTAP are needed, e.g. for the production of antibodies, vectors which direct high level expression of NTAP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of NTAP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH,

may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, supra; and Grant et al. (1987) *Methods Enzymol.* 153:516-54; Scorer, C. A. et al. (1994) *Bio/Technology* 5 12:181-184.)

Plant systems may also be used for expression of NTAP. Transcription of sequences encoding NTAP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV. (Takamatsu, N. (1987) *EMBO J.* 6:307-311.) Alternatively, plant promoters such as the small subunit of RUBISCO or 10 heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 15 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding NTAP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to 20 obtain infective virus which expresses NTAP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments 25 of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes.

For long term production of recombinant proteins in mammalian systems, stable expression of NTAP in cell lines is preferred. For example, sequences encoding NTAP can be 30 transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of 35 cells which successfully express the introduced sequences. Resistant clones of stably transformed

cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* or *ap<sup>r</sup>* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; and Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* or *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. 77:3567-3570; Colbere-Garapin, F. et al (1981) J. Mol. Biol. 150:1-14; and Murry, supra.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP) (Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -D-glucuronoside, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. et al. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding NTAP is inserted within a marker gene sequence, transformed cells containing sequences encoding NTAP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding NTAP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding NTAP and that express NTAP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of NTAP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on NTAP is preferred, but a

competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St Paul, MN, Section IV; Coligan, J. E. et al. (1997 and periodic supplements) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York, NY; and Maddox, D.E. et al. (1983) J. Exp. Med. 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding NTAP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding NTAP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison, WI), and U.S. Biochemical Corp. (Cleveland, OH). Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding NTAP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode NTAP may be designed to contain signal sequences which direct secretion of NTAP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to specify protein targeting, folding, and/or activity.

Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC, Manassas, VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding NTAP may be ligated to a heterologous sequence resulting in translation of a

fusion protein in any of the aforementioned host systems. For example, a chimeric NTAP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of NTAP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the NTAP encoding sequence and the heterologous protein sequence, so that NTAP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel, F. M. et al. (1995 and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch 10. A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled NTAP may be achieved in vitro using the TNT™ rabbit reticulocyte lysate or wheat germ extract systems (Promega, Madison, WI). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, preferably <sup>35</sup>S-methionine.

Fragments of NTAP may be produced not only by recombinant production, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the Applied Biosystems 431A Peptide Synthesizer (Perkin-Elmer Corp.). Various fragments of NTAP may be synthesized separately and then combined to produce the full length molecule.

## THERAPEUTICS

Partial chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of NTAP and various known neurotransmission associated proteins. In addition, NTAP is expressed in cancer and immortalized cell lines, and in inflammation and the immune response. Therefore, NTAP appears to play a role in cancer, and immune and neurological disorders.

Therefore, in one embodiment, NTAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a neurological disorder. Such disorders can include, but are not limited to, akathisia, Alzheimer's disease, amnesia, amyotrophic lateral sclerosis, bipolar disorder, catatonia, cerebral neoplasms, dementia, depression, diabetic neuropathy, Down's syndrome, tardive dyskinesia, dystonias, epilepsy, Huntington's disease, peripheral neuropathy, multiple sclerosis, neurofibromatosis, Parkinson's disease, paranoid psychoses, postherpetic neuralgia, schizophrenia, and Tourette's disorder.

In another embodiment, a vector capable of expressing NTAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a neurological disorder including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified NTAP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a neurological disorder including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of NTAP may be administered to a subject to treat or prevent a neurological disorder including, but not limited to, those listed above.

In a further embodiment, an antagonist of NTAP may be administered to a subject to treat or prevent a cancer. Such a cancer may include, but is not limited to, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. In one aspect, an antibody which specifically binds NTAP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express NTAP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding NTAP may be administered to a subject to treat or prevent a cancer including, but not limited to, those described above.

In a further embodiment, an antagonist of NTAP may be administered to a subject to treat or prevent an immune disorder. Such a disorder may include, but is not limited to, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia

with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's  
5 syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.

In an additional embodiment, a vector expressing the complement of the polynucleotide  
10 encoding NTAP may be administered to a subject to treat or prevent an immune disorder including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination  
15 therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of NTAP may be produced using methods which are generally known in the art. In particular, purified NTAP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind NTAP. Antibodies to NTAP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments,  
25 and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of polyclonal antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with NTAP or with any fragment or oligopeptide thereof which has immunogenic properties. Rats and mice are preferred hosts for  
30 downstream applications involving monoclonal antibody production. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and  
35 Corynebacterium parvum are especially preferable. (For review of methods for antibody

production and analysis, see, e.g., Harlow, E. and Lane, D. (1988) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.)

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to NTAP have an amino acid sequence consisting of at least about 5 amino acids, and, more preferably, of at least about 14 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of NTAP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to NTAP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci.* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci.* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce NTAP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton D.R. (1991) *Proc. Natl. Acad. Sci.* 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci.* 86:3833-3837; and Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for NTAP may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the

desired specificity and minimal cross-reactivity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between NTAP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering NTAP epitopes is preferred, but a competitive binding assay may also be employed. (Maddox, supra.)

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for NTAP. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of NTAP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple NTAP epitopes, represents the average affinity, or avidity, of the antibodies for NTAP. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular NTAP epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the NTAP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of NTAP, preferably in active form, from the antibody. (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, D. C.; and Liddell, J. E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York, NY.)

The titre and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is preferred for use in procedures requiring precipitation of NTAP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding NTAP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding NTAP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding NTAP. Thus, complementary molecules or fragments may be used to modulate NTAP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger

fragments can be designed from various locations along the coding or control regions of sequences encoding NTAP.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the  
5 targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding NTAP. (See, e.g., Sambrook, supra; and Ausubel, supra.)

Genes encoding NTAP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding NTAP. Such  
10 constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing  
15 complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding NTAP. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix  
20 pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY, pp. 163-177.) A complementary sequence or antisense molecule may also be  
25 designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For  
30 example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding NTAP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20  
ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may  
35 be evaluated for secondary structural features which may render the oligonucleotide inoperable.

The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These  
5 include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding NTAP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA,  
10 constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs  
15 and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally  
20 suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nature Biotechnology 15:462-466.)

25 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier,  
30 for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of NTAP, antibodies to NTAP, and mimetics, agonists, antagonists, or inhibitors of NTAP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions  
35 may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

5           In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

10           Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

15           Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as  
20 methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

          Dragee cores may be used in conjunction with suitable coatings, such as concentrated  
25 sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

          Pharmaceutical preparations which can be used orally include push-fit capsules made of  
30 gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

35           Pharmaceutical formulations suitable for parenteral administration may be formulated in

aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be  
5 prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

10 For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

15 The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acid. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2%  
20 sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of NTAP, such labeling would include amount, frequency, and method of administration.

25 Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in  
30 cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example  
35 NTAP or fragments thereof, antibodies of NTAP, and agonists, antagonists or inhibitors of NTAP,

which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The dose ratio of therapeutic to toxic effects is the therapeutic index, and it can be expressed as the  $ED_{50}/LD_{50}$  ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

25

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind NTAP may be used for the diagnosis of disorders characterized by expression of NTAP, or in assays to monitor patients being treated with NTAP or agonists, antagonists, or inhibitors of NTAP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for NTAP include methods which utilize the antibody and a label to detect NTAP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known

in the art and may be used.

A variety of protocols for measuring NTAP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of NTAP expression. Normal or standard values for NTAP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to NTAP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, preferably by photometric means. Quantities of NTAP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding NTAP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of NTAP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of NTAP, and to monitor regulation of NTAP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding NTAP or closely related molecules may be used to identify nucleic acid sequences which encode NTAP. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine whether the probe identifies only naturally occurring sequences encoding NTAP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably have at least 50% sequence identity to any of the NTAP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from a sequence selected from the group consisting of SEQ ID NO:7-12 or from genomic sequences including promoters, enhancers, and introns of the NTAP gene.

Means for producing specific hybridization probes for DNAs encoding NTAP include the cloning of polynucleotide sequences encoding NTAP or NTAP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as <sup>32</sup>P or <sup>35</sup>S, or by enzymatic labels,

such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding NTAP may be used for the diagnosis of a disorder associated with expression of NTAP. Examples of such a disorder include, but are not limited to, a neurological disorder such as akathisia, Alzheimer's disease, amnesia, amyotrophic lateral sclerosis, bipolar disorder, catatonia, cerebral neoplasms, dementia, depression, diabetic neuropathy, Down's syndrome, tardive dyskinesia, dystonias, epilepsy, Huntington's disease, peripheral neuropathy, multiple sclerosis, neurofibromatosis, Parkinson's disease, paranoid psychoses, postherpetic neuralgia, schizophrenia, and Tourette's disorder; a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an immune disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.. The polynucleotide sequences encoding NTAP may be used in Southern or Northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patients to detect altered NTAP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding NTAP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding NTAP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly

altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding NTAP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

5 In order to provide a basis for the diagnosis of a disorder associated with expression of NTAP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding NTAP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from  
10 normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated,  
15 hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of a relatively high amount of transcript in biopsied  
20 tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

25 Additional diagnostic uses for oligonucleotides designed from the sequences encoding NTAP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding NTAP, or a fragment of a polynucleotide complementary to the polynucleotide encoding NTAP, and will be employed under optimized conditions for  
30 identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of NTAP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol.  
35 Methods 159:235-244; and Duplaa, C. et al. (1993) Anal. Biochem. 229-236.) The speed of

quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding NTAP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, R.A. (ed.) Molecular Biology and Biotechnology, VCH Publishers New York, NY, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) site. Correlation between the location of the gene encoding NTAP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical

mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, NTAP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between NTAP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with NTAP, or fragments thereof, and washed. Bound NTAP is then detected by methods well known in the art. Purified NTAP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding NTAP specifically compete with a test compound for binding NTAP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with NTAP.

In additional embodiments, the nucleotide sequences which encode NTAP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/091,677, are hereby expressly incorporated by reference.

## EXAMPLES

### I. Construction of cDNA Libraries

RNA was purchased from CLONTECH Laboratories, Inc. or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIzol™ (Life Technologies, Inc.), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX™ latex particles (QIAGEN Inc., Valencia, CA), or an OLIGOTEX™ mRNA purification kit (QIAGEN Inc.). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE™ mRNA purification kit (Ambion, Austin, TX).

In some cases, Stratagene, Inc. (La Jolla, CA), was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP™ vector system (Stratagene, Inc.) or SUPERScript™ plasmid system (Life Technologies, Inc.), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, *supra*, 1997, units 5.1-6.6) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL® S1000, SEPHAROSE® CL2B, or SEPHAROSE® CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., pBLUESCRIPT® plasmid (Stratagene, Inc.), pSPORT™ 1 plasmid (Life Technologies, Inc.), or pINCY (Incyte Pharmaceuticals, Inc., Palo Alto, CA). Recombinant plasmids were transformed into competent *E. coli* cells, e.g., the XL1-Blue, XL1-BlueMRF, or SOLR™ strains (Stratagene, Inc.), or DH5α™, DH10B, or ElectroMAX DH10B competent cells (Life Technologies, Inc.).

### II. Isolation of cDNA Clones

Plasmids were recovered from host cells by *in vivo* excision, using the UNIZAP™ vector system (Stratagene, Inc.), or by cell lysis. Plasmids were purified using at least one of the

following: a Magic or WIZARD® Minipreps DNA purification system (Promega Corp.); an AGTC® Miniprep purification kit (Edge Biosystems, Gaithersburg, MD); the QIAWELL® 8 Plasmid, QIAWELL® 8 Plus Plasmid, or the QIAWELL® 8 Ultra Plasmid purification systems (QIAGEN Inc.); or the R.E.A.L.™ Prep 96 plasmid kit (QIAGEN Inc.). Following precipitation, 5 plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format. (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and 10 stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN® dye (Molecular Probes, Inc., Eugene, OR) and a Fluoroskan II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

### III. Sequencing and Analysis

The cDNAs were prepared for sequencing using either an ABI CATALYST 800 (Perkin- 15 Elmer ) or a MICROLAB® 2200 (Hamilton) sequencing preparation system in combination with Peltier PTC-200 thermal cyclers (MJ Research, Inc., Watertown, MA). The cDNAs were sequenced using the ABI PRISM™ 373 or 377 sequencing systems and ABI protocols, base calling software, and kits (Perkin-Elmer ). Alternatively, solutions and dyes from Amersham Pharmacia Biotech, Ltd. were used in place of the ABI kits. In some cases, reading frames were 20 determined using standard methods (Ausubel, supra). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA, extension, and shotgun sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the software programs used, 25 corresponding algorithms, references, and cutoff parameters used where applicable. The references cited in the third column of Table 5 are incorporated by reference herein. Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering Co.) and LASERGENE software (DNASTAR Inc.).

The polynucleotide sequences were validated by removing vector, linker, and polyA 30 sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS to acquire annotation, using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length 35 polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened

for open reading frames using programs based on GeneMark, BLAST, and FASTA. This was followed by translation of the full length polynucleotide sequences to derive the corresponding full length amino acid sequences. These full length polynucleotide and amino acid sequences were subsequently analyzed by querying against databases such as the GenBank databases described  
 5 above and SwissProt, BLOCKS, PRINTS, PFAM, and Prosite.

#### IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7;  
 10 and Ausubel, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ® database (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any  
 15 particular match is categorized as exact or similar.

The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the  
 20 length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of Northern analysis are reported as a list of libraries in which the transcript  
 25 encoding NTAP occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

#### V. Extension of NTAP Encoding Polynucleotides

30 Full-length nucleic acid sequences (SEQ ID NO:7-12) were produced by extension of the component fragments described in Table 1, Column 5, using oligonucleotide primers based on those fragments. For each nucleic acid sequence, one primer was synthesized to initiate extension of an antisense polynucleotide, and the other was synthesized to initiate extension of a sense polynucleotide. Primers were used to facilitate the extension of the known sequence "outward"  
 35 generating amplicons containing new unknown nucleotide sequence for the region of interest. The

initial primers were designed from the cDNA using OLIGO™ 4.06 (National Biosciences, Plymouth, MN), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension is necessary or desired, additional sets of primers are designed to further extend the known region.

High fidelity amplification was obtained by following the instructions for the XL-PCR™ kit (Perkin-Elmer Corp.) and thoroughly mixing the enzyme and reaction mix. PCR was performed using the PTC-200 thermal cycler (MJ Research, Inc.), beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, with the following parameters:

15	Step 1	94° C for 1 min (initial denaturation)
	Step 2	65° C for 1 min
	Step 3	68° C for 6 min
	Step 4	94° C for 15 sec
	Step 5	65° C for 1 min
	Step 6	68° C for 7 min
20	Step 7	Repeat steps 4 through 6 for an additional 15 cycles
	Step 8	94° C for 15 sec
	Step 9	65° C for 1 min
	Step 10	68° C for 7:15 min
	Step 11	Repeat steps 8 through 10 for an additional 12 cycles
25	Step 12	72° C for 8 min
	Step 13	4° C (and holding)

A 5 µl to 10 µl aliquot of the reaction mixture was analyzed by electrophoresis on a low concentration (about 0.6% to 0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were excised from the gel, purified using QIAQUICK™ kit (QIAGEN Inc.), and trimmed of overhangs using Klenow enzyme to facilitate religation and cloning.

After ethanol precipitation, the products were redissolved in 13 µl of ligation buffer, 1 µl T4-DNA ligase (15 units) and 1 µl T4 polynucleotide kinase were added, and the mixture was incubated at room temperature for 2 to 3 hours, or overnight at 16° C. Competent *E. coli* cells (in 40 µl of appropriate media) were transformed with 3 µl of ligation mixture and cultured in 80 µl of SOC medium. (See, e.g., Sambrook, *supra*, Appendix A, p. 2.) After incubation for one hour at 37°C, the *E. coli* mixture was plated on Luria Bertani (LB) agar (See, e.g., Sambrook, *supra*, Appendix A, p. 1) containing carbenicillin (2x carb). The following day, several colonies were

randomly picked from each plate and cultured in 150  $\mu$ l of liquid LB/2x carb medium placed in an individual well of an appropriate commercially-available sterile 96-well microtiter plate. The following day, 5  $\mu$ l of each overnight culture was transferred into a non-sterile 96-well plate and, after dilution 1:10 with water, 5  $\mu$ l from each sample was transferred into a PCR array.

- 5 For PCR amplification, 18  $\mu$ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction were added to each well. Amplification was performed using the following conditions:

	Step 1	94° C for 60 sec
10	Step 2	94° C for 20 sec
	Step 3	55° C for 30 sec
	Step 4	72° C for 90 sec
	Step 5	Repeat steps 2 through 4 for an additional 29 cycles
	Step 6	72° C for 180 sec
15	Step 7	4° C (and holding)

Aliquots of the PCR reactions were run on agarose gels together with molecular weight markers. The sizes of the PCR products were compared to the original partial cDNAs, and appropriate clones were selected, ligated into plasmid, and sequenced.

- 20 In like manner, the nucleotide sequence of SEQ ID NO:7-12 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for 5' extension, and an appropriate genomic library.

#### **VI. Labeling and Use of Individual Hybridization Probes**

- Hybridization probes derived from SEQ ID NO:7-12 are employed to screen cDNAs, 25 genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO™ 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase 30 (DuPont NEN®, Boston, MA). The labeled oligonucleotides are substantially purified using a Sephadex™ G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

- 35 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature

under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR™ film (Kodak, Rochester, NY) is exposed to the blots to film for several hours, hybridization patterns are compared visually.

## VII. Microarrays

- 5 A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements.
- 10 After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

- Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may
- 15 comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE™ software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g.,
- 20 UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; and Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

## VIII. Complementary Polynucleotides

- 25 Sequences complementary to the NTAP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring NTAP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO™ 4.06 software and the coding sequence of NTAP. To inhibit
- 30 transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the NTAP-encoding transcript.

## IX. Expression of NTAP

- Expression and purification of NTAP is achieved using bacterial or virus-based
- 35 expression systems. For expression of NTAP in bacteria, cDNA is subcloned into an appropriate

vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express NTAP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of NTAP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding NTAP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E. K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, NTAP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Pharmacia, Piscataway, NJ). Following purification, the GST moiety can be proteolytically cleaved from NTAP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak, Rochester, NY). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN Inc, Chatsworth, CA). Methods for protein expression and purification are discussed in Ausubel, F. M. et al. (1995 and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch 10, 16. Purified NTAP obtained by these methods can be used directly in the following activity assay.

#### **X. Demonstration of NTAP Activity**

NTAP, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton et al. (1973) Biochem. J. 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled NTAP, washed, and any wells with labeled NTAP complex are assayed. Data obtained using different concentrations of NTAP are used to calculate values for the number, affinity, and association of NTAP with the

candidate molecules.

#### **XI. Functional Assays**

NTAP function is assessed by expressing the sequences encoding NTAP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT™ plasmid (Life Technologies) and pCR™ 3.1 plasmid (Invitrogen, Carlsbad, CA) both of which contain the cytomegalovirus promoter. 5-10 μg of recombinant vector are transiently transfected into a human cell line, preferably of endothelial or hematopoietic origin, using either liposome formulations or electroporation. 1-2 μg of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP) (Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP, and to evaluate properties, for example, their apoptotic state. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M. G. (1994) Flow Cytometry, Oxford, New York, NY.

The influence of NTAP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding NTAP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success, NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding NTAP and other genes of interest can be analyzed by Northern analysis or microarray techniques.

**XII. Production of NTAP Specific Antibodies**

NTAP substantially purified using polyacrylamide gel electrophoresis (PAGE)(see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

5 Alternatively, the NTAP amino acid sequence is analyzed using LASERGENE™ software (DNASTAR Inc.) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel supra, ch. 11.)

10 Typically, oligopeptides 15 residues in length are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to KLH (Sigma Aldrich, St. Louis, MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for  
15 anti-peptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

**XIII. Purification of Naturally Occurring NTAP Using Specific Antibodies**

Naturally occurring or recombinant NTAP is substantially purified by immunoaffinity chromatography using antibodies specific for NTAP. An immunoaffinity column is constructed  
20 by covalently coupling anti-NTAP antibody to an activated chromatographic resin, such as CNBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing NTAP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of NTAP (e.g., high ionic strength  
25 buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/NTAP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and NTAP is collected.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of  
30 the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	7	238506	SINTNOT02	238506H1 (SINTNOT02), 881185R1 (THYRNOT02), 1809706F6 and 1809880T6 (PROSTUT12), 2371286F6 (ADRENOT07), 3558166H1 (LUNGNOT31)
2	8	414692	BRSTNOT01	030604F1 (THPINOB01), 061690F1 (LUNGNOT01), 414692H1 (BRSTNOT01), 1533952F1 (SPLNNOT04)
3	9	998868	KIDNTUT01	484199H1 and 484199R6 (HNT2RAT01), 489188H1 (HNT2AGT01), 998868H1 (KIDNTUT01), 1944742R6 (PITUNOT01), 2206131F6 (SPLNFET02), 3509110H1 (CONCNOT01)
4	10	1296451	PGANNOT03	000531R7 (U937NOT01), 623274R6 (PGANNOT01), 1295462F1 (PGANNOT03), 1296451H1 (PGANNOT03), 1664686F6 (BRSTNOT09), 2824843H1 (ADRETUT06)
5	11	1739035	COLNNOT22	1469082F6 and 1469082T1 (PANCTUT02), 1665101H1 (BRSTNOT09), 1739035F6 and 1739035H1 (COLNNOT22), 1998944R6 (BRSTTUT03), 3770403H1 (BRSTNOT25)
6	12	2799056	NPOLNOT01	875877R1 (LUNGAST01), 1509734F1 (LUNGNOT14), 1979922R6, 1979922T6 and 1981971H1 (LUNGUT03), 2799056H1 (NPOLNOT01), SAWA01057F1

Table 2

Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Identification	Analytical Methods
1	251	S88 S174 S191 T205 S25 T199			neuronal protein	BLAST
2	238	S11 S12 T105 T117 S145 S206 S61		S21-Q45	NMDA receptor	BLAST PRINTS
3	408	T8 S33 T220 S225 T289 S330 S331 S58 S200 T284 Y165 Y372	N31 N204		Benzodiazepine receptor- associated protein	BLAST
4	272	T2 S103 T126 T167 S177 S244		C43-C54 C74-C85 C139-C150	Putative mammalian homeotic protein	BLAST MOTIFS PRINTS BLOCKS
5	363	T192 T19 S65 S86 T213 T18 T47 S133 T159 T215 S253	N4 N14 N131 N251	P229-P240	rhodopsin	PRINTS
6	484	S102 T127 S298 T390 T140 S164 S404 Y236	N48 N264 N401		Putative odorant- binding protein	BLAST

Table 3

Seq ID NO:	Tissue Expression (Fraction of Total)	Disease Class (Fraction of Total)	Vector
7	Reproductive (0.275) Hematopoietic/Immune (0.200) Gastrointestinal (0.175)	Cancer (0.400) Inflammation (0.375) Fetal (0.175)	pBluescript
8	Reproductive (0.272) Cardiovascular (0.140) Hematopoietic/Immune (0.132)	Cancer (0.439) Inflammation (0.281) Fetal (0.140)	pBluescript
9	Cardiovascular (0.200) Nervous (0.200) Developmental (0.133)	Fetal (0.467) Cancer (0.400) Inflammation (0.200)	pSPORT1
10	Nervous (0.278) Developmental (0.250) Reproductive (0.222)	Cancer (0.333) Fetal (0.333) Inflammation (0.167)	pINCY
11	Reproductive (0.381) Gastrointestinal (0.286) Developmental (0.119)	Cancer (0.548) Inflammation (0.261) Fetal (0.214)	pINCY
12	Cardiovascular (0.444) Reproductive (0.278) Developmental (0.167)	Cancer (0.667) Fetal (0.167) Inflammation (0.167)	pINCY

Table 4

Nucleotide SEQ ID NO:	Clone ID	Library	Library Comment
7	238506	SINTNOT02	Library was constructed using RNA isolated from the small intestine of a 55-year-old Caucasian female, who died from a subarachnoid hemorrhage. Serologies were positive for cytomegalovirus (CMV).
8	414692	BRSTNOT01	Library was constructed using RNA isolated from the breast tissue of a 56-year-old Caucasian female who died in a motor vehicle accident.
9	998868	KIDNTUT01	Library was constructed using RNA isolated from the kidney tumor tissue removed from an 8-month-old female during nephroureterectomy. Pathology indicated Wilms' tumor (nephroblastoma), which involved 90 percent of the renal parenchyma.
10	1296451	PGANNOT03	Library was constructed using RNA isolated from paraganglionic tumor tissue removed from the intra-abdominal region of a 46-year-old Caucasian male during exploratory laparotomy. Pathology indicated a benign paraganglioma and was associated with a grade 2 renal cell carcinoma, clear cell type, which did not penetrate the capsule.
11	1739035	COLNNOT22	Library was constructed using RNA isolated from colon tissue removed from a 56-year-old Caucasian female with Crohn's disease during a partial resection of the small intestine. Pathology indicated Crohn's disease of the ileum and ileal-colonic anastomosis, causing a fistula at the anastomotic site that extended into pericolonic fat. The ileal mucosa showed linear and punctate ulcers with intervening normal tissue.
12	2799056	NPOLNOT01	Library was constructed using RNA isolated from nasal polyp tissue removed from a 78-year-old Caucasian male during a nasal polypectomy. Pathology indicated a nasal polyp and striking eosinophilia. Patient history included asthma and nasal polyps.

TABLE 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABU/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA includes five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: z-score = 10 or greater Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS and PRINTS databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and Probability value= 1.0E-3 or less
PFAM	A Hidden Markov Models-based application useful for protein family search.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits, depending on individual protein families

TABLE 5 (con)

Program	Description	Reference	Parameter Threshold
GeneMark	A gene prediction algorithm that is based on inhomogeneous Markov chain models and is useful for DNA sequence analysis and particularly for gene prediction.	Borodovsky, M. and J. McIninch (1993) Computers Chem. 17: 123-133; Blattner, F.R. et al. (1993) Nucleic Acids Res. 21: 5408-5417.	Score= 0.4 or greater
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribnikov, M. et al. (1988) CABIOS 4:61-66; Gribnikov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Score= 4.0 or greater
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147: 195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6, and fragments thereof.
2. A substantially purified variant having at least 90% amino acid identity to the amino acid sequence of claim 1.
3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.
4. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 3.
5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.
6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide sequence of claim 3.
7. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12, and fragments thereof.
8. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 7.
9. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 7.
10. An expression vector comprising at least a fragment of the polynucleotide of claim 3.
11. A host cell comprising the expression vector of claim 10.

12. A method for producing a polypeptide, the method comprising the steps of:
- a) culturing the host cell of claim 11 under conditions suitable for the expression of the polypeptide; and
  - b) recovering the polypeptide from the host cell culture.
- 5
13. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.
14. A purified antibody which specifically binds to the polypeptide of claim 1.
- 10
15. A purified agonist of the polypeptide of claim 1.
16. A purified antagonist of the polypeptide of claim 1.
- 15
17. A method for treating or preventing neurological disorder, the method comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 13.
18. A method for treating or preventing a cancer, the method comprising
- 20 administering to a subject in need of such treatment an effective amount of the antagonist of claim 16.
19. A method for treating or preventing an immune disorder, the method comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim
- 25 16.
20. A method for detecting a polynucleotide, the method comprising the steps of:
- (a) hybridizing the polynucleotide of claim 6 to at least one of the nucleic acids in a biological sample, thereby forming a hybridization complex; and
  - (b) detecting the hybridization complex, wherein the presence of the
- 30 hybridization complex correlates with the presence of the polynucleotide encoding the polypeptide in the biological sample.
21. The method of claim 20 further comprising amplifying the polynucleotide prior to
- 35 hybridization.

## SEQUENCE LISTING

&lt;110&gt; INCYTE PHARMACEUTICALS, INC.

LAL, Preeti  
 TANG, Y. Tom  
 YUE, Henry  
 CORLEY, Neil C.  
 GUEGLER, Karl J.  
 GORGONE, Gina A.  
 BAUGHN, Mariah R.  
 PATTERSON, Chandra

&lt;120&gt; NEUROTRANSMISSION ASSOCIATED PROTEINS

&lt;130&gt; PF-0551 PCT

&lt;140&gt; To Be Assigned

&lt;141&gt; Herewith

&lt;150&gt; 60/091,667

&lt;151&gt; 1998-07-02

&lt;160&gt; 12

&lt;170&gt; PERL Program

&lt;210&gt; 1

&lt;211&gt; 251

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 238506

&lt;400&gt; 1

Leu	Leu	Lys	Pro	Gly	Leu	Arg	Ala	Val	Val	Gly	Gly	Ala	Ala	Ala	1	5	10	15
Val	Ser	Thr	Gln	Ala	Met	His	Asn	Gly	Ser	Pro	Lys	Ser	Ser	Ala	20	25	30	
Ser	Gln	Ala	Gly	Ala	Ala	Ala	Gly	Gln	Gly	Ala	Pro	Ala	Pro	Ala	35	40	45	
Pro	Ala	Ser	Gln	Glu	Pro	Leu	Pro	Ile	Ala	Gly	Pro	Ala	Thr	Ala	50	55	60	
Pro	Ala	Pro	Arg	Pro	Leu	Gly	Ser	Ile	Gln	Arg	Pro	Asn	Ser	Phe	65	70	75	
Leu	Phe	Arg	Ser	Ser	Ser	Gln	Ser	Gly	Ser	Gly	Pro	Ser	Ser	Pro	80	85	90	
Asp	Ser	Val	Leu	Arg	Pro	Arg	Arg	Tyr	Pro	Gln	Val	Pro	Asp	Glu	95	100	105	
Lys	Asp	Leu	Met	Thr	Gln	Leu	Arg	Gln	Val	Leu	Glu	Ser	Arg	Leu	110	115	120	
Gln	Arg	Pro	Leu	Pro	Glu	Asp	Leu	Ala	Glu	Ala	Leu	Ala	Ser	Gly	125	130	135	
Val	Ile	Leu	Cys	Gln	Leu	Ala	Asn	Gln	Leu	Arg	Pro	Arg	Ser	Val	140	145	150	

Pro	Phe	Ile	His	Val	Pro	Ser	Pro	Ala	Val	Pro	Lys	Leu	Ser	Ala
				155					160					165
Leu	Lys	Ala	Arg	Lys	Asn	Val	Glu	Ser	Phe	Leu	Glu	Ala	Cys	Arg
				170					175					180
Lys	Met	Gly	Val	Pro	Glu	Ala	Asp	Leu	Cys	Ser	Pro	Ser	Asp	Leu
				185					190					195
Leu	Gln	Gly	Thr	Ala	Arg	Gly	Leu	Arg	Thr	Ala	Leu	Glu	Ala	Val
				200					205					210
Lys	Arg	Val	Gly	Gly	Lys	Ala	Leu	Pro	Pro	Leu	Trp	Pro	Pro	Ser
				215					220					225
Gly	Leu	Gly	Gly	Phe	Val	Val	Phe	Tyr	Val	Val	Leu	Met	Leu	Leu
				230					235					240
Leu	Tyr	Val	Thr	Tyr	Thr	Arg	Leu	Leu	Gly	Ser				
				245					250					

&lt;210&gt; 2

&lt;211&gt; 238

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 414692

&lt;400&gt; 2

Met	Ala	Asp	Pro	Asp	Pro	Arg	Tyr	Pro	Arg	Ser	Ser	Ile	Glu	Asp
1				5					10					15
Asp	Phe	Asn	Tyr	Gly	Ser	Ser	Val	Ala	Ser	Ala	Thr	Val	His	Ile
				20					25					30
Arg	Met	Ala	Phe	Leu	Arg	Lys	Val	Tyr	Ser	Ile	Leu	Ser	Leu	Gln
				35					40					45
Val	Leu	Leu	Thr	Thr	Val	Thr	Ser	Thr	Val	Phe	Leu	Tyr	Phe	Glu
				50					55					60
Ser	Val	Arg	Thr	Phe	Val	His	Glu	Ser	Pro	Ala	Leu	Ile	Leu	Leu
				65					70					75
Phe	Ala	Leu	Gly	Ser	Leu	Gly	Leu	Ile	Phe	Ala	Leu	Thr	Leu	Asn
				80					85					90
Arg	His	Lys	Tyr	Pro	Leu	Asn	Leu	Tyr	Leu	Leu	Phe	Gly	Phe	Thr
				95					100					105
Leu	Leu	Glu	Ala	Leu	Thr	Val	Ala	Val	Val	Val	Thr	Phe	Tyr	Asp
				110					115					120
Val	Tyr	Ile	Ile	Leu	Gln	Ala	Phe	Ile	Leu	Thr	Thr	Thr	Val	Phe
				125					130					135
Phe	Gly	Leu	Thr	Val	Tyr	Thr	Leu	Gln	Ser	Lys	Lys	Asp	Phe	Ser
				140					145					150
Lys	Phe	Gly	Ala	Gly	Leu	Phe	Ala	Leu	Leu	Trp	Ile	Leu	Cys	Leu
				155					160					165
Ser	Gly	Phe	Leu	Lys	Phe	Phe	Phe	Tyr	Ser	Glu	Ile	Met	Glu	Leu
				170					175					180
Val	Leu	Ala	Ala	Ala	Gly	Ala	Leu	Leu	Phe	Cys	Gly	Phe	Ile	Ile
				185					190					195
Tyr	Asp	Thr	His	Ser	Leu	Met	His	Lys	Leu	Ser	Pro	Glu	Glu	Tyr
				200					205					210
Val	Leu	Ala	Ala	Ile	Ser	Leu	Tyr	Leu	Asp	Ile	Ile	Asn	Leu	Phe

	215		220	225
Leu His Leu Leu Arg Phe Leu Glu Ala Val Asn Lys Lys				
	230		235	

<210> 3  
 <211> 408  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte Clone No: 998868

<400> 3  
 Met Gly Pro Tyr Asn Pro Asp Thr Cys Pro Glu Val Gly Phe Phe  
 1 5 10 15  
 Asp Val Leu Gly Asn Asp Arg Arg Arg Glu Trp Ala Ala Leu Gly  
 20 25 30  
 Asn Met Ser Lys Glu Asp Ala Met Val Glu Phe Val Lys Leu Leu  
 35 40 45  
 Asn Arg Cys Cys His Leu Phe Ser Thr Tyr Val Ala Ser His Lys  
 50 55 60  
 Ile Glu Lys Glu Glu Gln Asp Lys Lys Arg Gln Glu Glu Glu Glu  
 65 70 75  
 Arg Arg Arg Arg Glu Glu Glu Glu Arg Glu Arg Leu Pro Lys Glu  
 80 85 90  
 Glu Glu Lys Arg Arg Arg Glu Glu Glu Glu Arg Leu Arg Arg Ala  
 95 100 105  
 Ala Glu Glu Arg Arg Arg Ile Glu Glu Glu Arg Leu Arg Leu Glu  
 110 115 120  
 Gln Gln Lys Gln Gln Ile Met Ala Ala Leu Asn Ser Gln Thr Ala  
 125 130 135  
 Val Gln Phe Gln Gln Tyr Ala Ala Gln Gln Tyr Pro Gly Asn Tyr  
 140 145 150  
 Glu Gln Gln Gln Ile Leu Ile Arg Gln Leu Gln Glu Gln His Tyr  
 155 160 165  
 Gln Gln Tyr Met Gln Gln Leu Tyr Gln Val Gln Leu Ala Gln Gln  
 170 175 180  
 Gln Ala Ala Leu Gln Lys Gln Gln Glu Val Val Val Ala Gly Ser  
 185 190 195  
 Ser Leu Pro Thr Ser Ser Lys Val Asn Ala Thr Val Pro Ser Asn  
 200 205 210  
 Met Met Ser Val Asn Gly Gln Ala Lys Thr His Thr Asp Ser Ser  
 215 220 225  
 Glu Lys Glu Leu Glu Pro Glu Ala Ala Glu Glu Ala Leu Glu Asn  
 230 235 240  
 Gly Pro Lys Glu Ser Leu Pro Val Ile Ala Ala Pro Ser Met Trp  
 245 250 255  
 Thr Arg Pro Gln Ile Lys Asp Phe Lys Glu Lys Ile Gln Gln Asp  
 260 265 270  
 Ala Asp Ser Val Ile Thr Val Gly Arg Gly Glu Val Val Thr Val  
 275 280 285  
 Arg Val Pro Thr His Glu Glu Gly Ser Tyr Leu Phe Trp Glu Phe  
 290 295 300

PCT/US99/15121

```
<210> 4
<211> 272
<212> PRT
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<223> Incyte Clone No: 1296451
```

4/11

Val Gly Ile Val	Phe Leu Asn Lys Cys Glu Thr Trp Val Ser Asn	
	215	220 225
Leu Arg Tyr Asn His Met Leu Arg Lys Lys Lys Asn Leu Leu Leu		
	230	235 240
Gln Tyr Asn Ser Gly Glu Asp Leu Ala Val Asn Ile Ile Phe Pro		
	245	250 255
Glu Lys Ile Asp Met Thr Thr Phe Ser Lys Glu Ala Gly Asp Glu		
	260	265 270
Glu Ile		

&lt;210&gt; 5

&lt;211&gt; 363

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 1739035

&lt;400&gt; 5

Met Cys Leu Asn His Ser Asn Gln Phe Thr Gln Leu Gly Asn Ile	
1	5 10 15
Thr Glu Thr Thr Lys Phe Glu Lys Leu Ala Glu Asp Cys Lys Arg	
	20 25 30
Ser Met Asp Ile Leu Lys Gln Ala Phe Val Arg Gly Leu Pro Thr	
	35 40 45
Pro Thr Ala Arg Phe Glu Gln Arg Thr Phe Ser Val Ile Lys Ile	
	50 55 60
Phe Pro Asp Leu Ser Ser Asn Asp Met Leu Leu Phe Ile Val Lys	
	65 70 75
Gly Ile Asn Leu Pro Thr Pro Pro Gly Leu Ser Pro Gly Asp Leu	
	80 85 90
Asp Val Phe Val Arg Phe Asp Phe Pro Tyr Pro Asn Val Glu Glu	
	95 100 105
Ala Gln Lys Asp Lys Thr Ser Val Ile Lys Asn Thr Asp Ser Pro	
	110 115 120
Glu Phe Lys Glu Gln Phe Lys Leu Cys Ile Asn Arg Ser His Arg	
	125 130 135
Gly Phe Arg Arg Ala Ile Gln Thr Lys Gly Ile Lys Phe Glu Val	
	140 145 150
Val His Lys Gly Gly Leu Phe Lys Thr Asp Arg Val Leu Gly Thr	
	155 160 165
Ala Gln Leu Lys Leu Asp Ala Leu Glu Ile Ala Cys Glu Val Arg	
	170 175 180
Glu Ile Leu Glu Val Leu Asp Gly Arg Arg Pro Thr Gly Gly Arg	
	185 190 195
Leu Glu Val Met Val Arg Ile Arg Glu Pro Leu Thr Ala Gln Gln	
	200 205 210
Leu Glu Thr Thr Thr Glu Arg Trp Leu Val Ile Asp Pro Val Pro	
	215 220 225
Ala Ala Val Pro Thr Gln Val Ala Gly Pro Lys Gly Lys Ala Pro	
	230 235 240
Pro Val Pro Ala Pro Ala Arg Glu Ser Gly Asn Arg Ser Ala Arg	
	245 250 255

```

Pro Leu His Ser Leu Ser Val Leu Ala Phe Asp Gln Glu Arg Leu
      260                      265                      270
Glu Arg Lys Ile Leu Ala Leu Arg Gln Ala Arg Arg Pro Val Pro
      275                      280                      285
Pro Glu Val Ala Gln Gln Tyr Gln Asp Ile Met Gln Arg Ser Gln
      290                      295                      300
Trp Gln Arg Ala Gln Leu Glu Gln Gly Gly Val Gly Ile Arg Arg
      305                      310                      315
Glu Tyr Ala Ala Gln Leu Glu Arg Gln Leu Gln Phe Tyr Thr Glu
      320                      325                      330
Ala Ala Arg Arg Leu Gly Asn Asp Gly Ser Arg Asp Ala Ala Lys
      335                      340                      345
Glu Ala Leu Tyr Arg Arg Asn Leu Val Glu Ser Glu Leu Gln Arg
      350                      355                      360
Leu Arg Arg

```

&lt;210&gt; 6

&lt;211&gt; 484

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 2799056

&lt;400&gt; 6

```

Met Ala Gly Pro Trp Thr Phe Thr Leu Leu Cys Gly Leu Leu Ala
  1                      5                      10                      15
Ala Thr Leu Ile Gln Ala Thr Leu Ser Pro Thr Ala Val Leu Ile
      20                      25                      30
Leu Gly Pro Lys Val Ile Lys Glu Lys Leu Thr Gln Glu Leu Lys
      35                      40                      45
Asp His Asn Ala Thr Ser Ile Leu Gln Gln Leu Pro Leu Leu Ser
      50                      55                      60
Ala Met Arg Glu Lys Pro Ala Gly Gly Ile Pro Val Leu Gly Ser
      65                      70                      75
Leu Val Asn Thr Val Leu Lys His Ile Ile Trp Leu Lys Val Ile
      80                      85                      90
Thr Ala Asn Ile Leu Gln Leu Gln Val Lys Pro Ser Ala Asn Asp
      95                      100                     105
Gln Glu Leu Leu Val Lys Ile Pro Leu Asp Met Val Ala Gly Phe
      110                     115                     120
Asn Thr Pro Leu Val Lys Thr Ile Val Glu Phe His Met Thr Thr
      125                     130                     135
Glu Ala Gln Ala Thr Ile Arg Met Asp Thr Ser Ala Ser Gly Pro
      140                     145                     150
Thr Arg Leu Val Leu Ser Asp Cys Ala Thr Ser His Gly Ser Leu
      155                     160                     165
Arg Ile Gln Leu Leu His Lys Leu Ser Phe Leu Val Asn Ala Leu
      170                     175                     180
Ala Lys Gln Val Met Asn Leu Leu Val Pro Ser Leu Pro Asn Leu
      185                     190                     195
Val Lys Asn Gln Leu Cys Pro Val Ile Glu Ala Ser Phe Asn Gly
      200                     205                     210

```

Met Tyr Ala Asp	Leu Leu Gln Leu Val	Lys Val Pro Ile Ser Leu	
	215	220	225
Ser Ile Asp Arg	Leu Glu Phe Asp Leu	Leu Tyr Pro Ala Ile Lys	
	230	235	240
Gly Asp Thr Ile	Gln Leu Tyr Leu Gly	Ala Lys Leu Leu Asp Ser	
	245	250	255
Gln Gly Lys Val	Thr Lys Trp Phe Asn	Asn Ser Ala Ala Ser Leu	
	260	265	270
Thr Met Pro Thr	Leu Asp Asn Ile Pro	Phe Ser Leu Ile Val Ser	
	275	280	285
Gln Asp Val Val	Lys Ala Ala Val Ala	Ala Val Leu Ser Pro Glu	
	290	295	300
Glu Phe Met Val	Leu Leu Asp Ser Val	Leu Pro Glu Ser Ala His	
	305	310	315
Arg Leu Lys Ser	Ser Ile Gly Leu Ile	Asn Glu Lys Ala Ala Asp	
	320	325	330
Lys Leu Gly Ser	Thr Gln Ile Val Lys	Ile Leu Thr Gln Asp Thr	
	335	340	345
Pro Glu Phe Phe	Ile Asp Gln Gly His	Ala Lys Val Ala Gln Leu	
	350	355	360
Ile Val Leu Glu	Val Phe Pro Ser Ser	Glu Ala Leu Arg Pro Leu	
	365	370	375
Phe Thr Leu Gly	Ile Glu Ala Ser Ser	Glu Ala Gln Phe Tyr Thr	
	380	385	390
Lys Gly Asp Gln	Leu Ile Leu Asn Leu	Asn Asn Ile Ser Ser Asp	
	395	400	405
Arg Ile Gln Leu	Met Asn Ser Gly Ile	Gly Trp Phe Gln Pro Asp	
	410	415	420
Val Leu Lys Asn	Ile Ile Thr Glu Ile	Ile His Ser Ile Leu Leu	
	425	430	435
Pro Asn Gln Asn	Gly Lys Leu Arg Ser	Gly Val Pro Val Ser Leu	
	440	445	450
Val Lys Ala Leu	Gly Phe Glu Ala Ala	Glu Ser Ser Leu Thr Lys	
	455	460	465
Asp Ala Leu Val	Leu Thr Pro Ala Ser	Leu Trp Lys Pro Ser Ser	
	470	475	480
Pro Val Ser Gln			

&lt;210&gt; 7

&lt;211&gt; 1638

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 238506

&lt;400&gt; 7

```

tcggtagatg gtgggctgga ctcaggcttc cacagcggtg atagtggcag caagaggtgg 60
tctggaaatg agtcaacaga tgaattttca gagctgtcat tccggatctc agagctggcc 120
cgggagcccc ggggaccag agaacgcaag gaggatggct cagcggacgg agaccctgtg 180
cagattgact tcatcgacag ccatgtcccc ggggaggatg aagagcgagg cactgtggag 240
gagcagcgac caccgaatt aagccctggg gcaggggaca gggagagggc accaagcagc 300
aggcgggagg agccggcagg ggaggagcgg cggcgcccg acaccttgca gctgtggcag 360

```

```

gagcggggaac ggcggcagca gcagcagagc ggggctgtgg gggccccgag gaaggatagc 420
ctcttgaagc cagggctcag ggctgttgtg ggaggggccc cgcctgtgtc cactcaagcc 480
atgcacaacg gctcgccctaa gtccagtgcc tcccaagcag gggctgcagc ggggcaggga 540
gcccccgccc ctgcccctgc ctcccaagag ccccttccca tagctggacc agcgacagca 600
cctgtctccac ggccacttgg ctccattcag agaccaaaca gcttctctt cegttctctc 660
tctcagagtg gctcaggccc ttctcacca gactctgtcc tgagacctcg gcggtacccc 720
caggttccag atgagaagga cttaatgact cagctgcgcc aggtccttga gtcccggctg 780
cagcgggccc tgctgagga cctggccgag gctctggcca gtggggcat cctgtgccag 840
ctggccaacc agctacggcc gcgtccgtg cccttcatcc atgtgccctc ccctgtgtg 900
ccaaaactca gtgccctcaa ggctcggaag aatgtggaga gttttctaga agcctgtcga 960
aaaatggggg tgcctgaggg tgacctgtgc tcgccctcgg atctctcca gggcactgcc 1020
cgggggctgc ggaccgcgct ggaggccgtg aagcgggtgg ggggcaaggc cctaccgcc 1080
ctctggcccc cctctggctt gggcggttc gtctcttct acgtggtcct catgtgtgtg 1140
ctctatgtca cctacactcg gctcctgggt tcctaggccc caaaatcggc cctccctcac 1200
ccctttccct tcctctctat ttataaggtc cctgtccac cgcacccac ctgcggtgcc 1260
ttcagcccca accaaagaca ctagtgcacc cccttcacag aactgacct cagaggcccc 1320
actctgggtc cccagacccc tgggccccca gcctctggcc tccctccagt agccccacga 1380
gtccccacct ctcagtgtg acggtgcctt catgtcccc cgggccctgc cctgcctc 1440
tgtaccccg tgggggtggc aggagctgga gtctcccc tctctctgtg cctccctt 1500
cccccccaa cagctgtat gggggggcta aattatctt atttttaga gaggatctat 1560
atgtgtagg gttcgggggc caggccgggt ccctatctt gtgtataaac tgtacagacc 1620
gtgaaaagaa aaaaaaaa 1638

```

&lt;210&gt; 8

&lt;211&gt; 1015

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 414692

&lt;400&gt; 8

```

cgctttctcc gccagctgg aatttttgaa gcgagaaaat cgactcgctc ggtgttcgcc 60
cgccgacgcc gcacggcttg ctggggctgg gctcttctc gcggaagtgg ggaggaggcg 120
gttgcggtta gtggaccggg accggtaggg gtgctgttg catcatggct gaccccgacc 180
ccgggtaccc tcgtcctcgt atcgaggacg acttcaacta tggcagcagc gtggcctccg 240
ccaccgtgca catccgaatg gcctttctga gaaaagtcta cagcattctt tctctgcagg 300
ttctcttaac tacagtgaact tcaacagttt ttttatactt tgagtctgta cggacatttg 360
tacatgagag tctgcctta attttgctgt ttgcctcgg atctctgggt ttgatttttg 420
cgttgacttt aaacagacat aagtatcccc ttaacctgta cctactttt ggatttacgc 480
tgttggaagc tctgactgtg gcagttgttg ttactttcta tgatgtatat attattctgc 540
aagctttcat actgactact acagtatttt ttggtttgac tgtgtatact ctacaatcta 600
agaaggattt cagcaaattt ggagcagggc tgtttgctct tttgtggata ttgtgcctgt 660
caggattctt gaagtttttt ttttatagt agataatgga gttggtctta gccgctgcag 720
gagcccttct tttctgtgga ttcacatct atgacacaca ctactgatg cataaactgt 780
cacctgaaga gtacgtatta gctgccatca gcctctact ggatatcatc aatctattcc 840
tgcacctgtt acggtttctg gaagcagtta ataaaaagta attaaaagta tctcagctca 900
actgaagaac aacaaaaaaa atttaacgag aaaaaaggat taaagtaatt ggaagcagta 960
tatagaaact gtttcattaa gtaataaagt ttgaaacaat gattaaaaaa aaaaa 1015

```

&lt;210&gt; 9

&lt;211&gt; 1481

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 998868

&lt;400&gt; 9

```

gcggcggtcg gagcagcgct ggggttttcgg cctggaggag ttgtacggcc tggcactgcg 60
cttcttcaaa gaaaaagatg gcaaagcatt tcatccaact tatgaagaaa aattgaagct 120
tgtggcactg cataagcaag ttcttatggg cccatataat ccagacactt gtcctgaggt 180
tggattcttt gatgtgttgg ggaatgacag gaggagagaa tgggcagccc tgggaaacat 240
gtctaaagag gatgccatgg tggagtttgt caagctctta aatagggtgt gccatctctt 300
ttcaacatat gttgcgtccc acaaaataga gaaggaagag caagacaaaa aaaggcagga 360
ggaagaggag cgaaggcggc gtgaagagga agaaagagaa cgtctgccaa aggaggaaga 420
gaaacgtagg agagaagaag aggaaaggct tcgacgggag gcagaggaaa ggagacggat 480
agaagaagaa aggtctcggt tggagcagca aaagcagcag ataatggcag ctttaaaactc 540
ccagactgcc gtgcagtcc agcagtatgc agcccaacag tatccaggga actacgaaca 600
gcagcaaatt ctcatccgcc agttgcagga gcaacactat cagcagtaca tgcagcagtt 660
gtatcaagtc cagcttgccac agcaacaggc agcattacag aaacaacagg aagtagtagt 720
ggctgggtct tccttgccca catcatcaaa agtgaatgca actgtacca gtaatatgat 780
gtcagttaat ggacaggcca aaacacacac tgacagctcc gaaaaagaac tgggaaccaga 840
agctgcagaa gaagccctgg agaattggacc aaaagaatct cttccagtaa tagcagctcc 900
atccatgtgg acacgacctc agatcaaaga cttcaaagag aagattcagc aggatgcaga 960
ttccgtgatt acagtggggc gaggagaagt ggtcactggt cgagtaccca cccatgaaga 1020
aggatcatat ctcttttggg aatttgccac agacaattat gacattgggt ttgggggtga 1080
ttttgaatgg acagactctc caaacactgc tgtcagcggt catgtcagtg agtccagcga 1140
tgacgacgag gaggaagaag aaaacatcgg ttgtgaagag aaagccaaaa agaattgcaa 1200
caagcctttg ctggatgaga ttgtgcctgt gtaccgacgg gactgtcatg aggaggtgta 1260
tgctggcagc catcaatatc caggagagag agtctatctc ctcaagtttg acaactccta 1320
ctctttgttg cggtcaaaaat cagtctacta cagagtctat tatactagat aaaaatgttg 1380
ttacaaagtc tggagtctag ggttgggcag aagatgacat ttaatttggg gatttctttt 1440
tacttttgtg gagcattaga gtcacagttt accttattga t 1481

```

&lt;210&gt; 10

&lt;211&gt; 1212

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 1296451

&lt;400&gt; 10

```

cgcgcacgag cagcccgggt cagccctggc tttcccctcg ctgcgcgccc gcgccccctt 60
tcgctgccgc aaccagaagc ccagtgcggc gccaggagcc ggacccgcgc ccgcaccgct 120
cccgggaccg cgaccccggc cgcccagaga tgaccgcgac cgaagccctc ctgcgcgtcc 180
tcttgctcct gctggctttc ggccacagca cctatggggc tgaatgcttc cgggcctgca 240
acccccaaaa tggattctgc gaggatgaca atgtttgcag gtgccagcct ggctggcagg 300
gtcccccttg tgaccagtgc gtgacctctc ccggtgcct tcacggactc tgtggagaac 360
ccgggcagtg catttgacc gacggctggg acggggagct ctgtgataga gatgttcggg 420
cctgtctctc ggccccctgt gccaaacaac ggtactcggg aaaggactgc cagaaaaagg 480
acgggccctg tgtgatcaac ggctccccct gccagcacgg aggcacctgc gtggatgatg 540
agggccgggc ctcccatgcc tctgctgtgt gcccccttg cttctcaggc aatttctgag 600
agatcgtggc cagcccgctg cagaacgggg gcacctgcct gcagcacacc cagccggagc 660
accgcctcct gaaggtgtcc atgaaagagc tcaacaagaa aacccctctc ctcccgagg 720
gccaggccat ctgcttcacc atcctggggc tgcctaccag cctgggtggg ctgggcactg 780
tgggtatcgt cttcctcaac aagtgcgaga cctgggtgtc caacctgcgc tacaaccaca 840
tgctgcggaa gaagaagaac ctgctgcttc agtacaacag cggggaggag ctggccgtca 900

```

```

acatcatctt ccccgagaag atcgacatga ccaccttcag caaggaggcc ggcgacgagg 960
agatctaagc agcgttccca cagccccctc tagattcttg gagttccgca gagcttacta 1020
taacgcggtct gtccctaactt ttgtggtggt cgctatctct tgtgtcaaata ctggtgaacg 1080
ctacgcttac atatattgtc tttgtgctgc tgtgtgacaa acgcaatgca aaaacaatcc 1140
tctttctctc tcttaatgca tgatacagaa taataataag aatttcatct ttaaatgaga 1200
tctggaattt ta 1212

```

&lt;210&gt; 11

&lt;211&gt; 1658

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 1739035

&lt;400&gt; 11

```

tggccccgggt ctgtctcagg aggccgcccc gcgctatggt gaactcacca agctcatacg 60
gcagcagcac gagatgtgcc tgaaccactc aaaccaattc acccagctgg gcaacatcac 120
tgaaaccacc aagtttgaaa agttggcgga ggactgtaag cggagcatgg acattctgaa 180
gcaagccttc gtccgggggtc tccccacgcc caccgccccg tttgagcaaa ggaccttcag 240
cgatcatcaag atcttccctg acctcagcag caacgacatg ctctcttca tctgaaggg 300
catcaacttg cccacacccc caggactgtc ccctggcgat ctggatgtct ttgttcggtt 360
tgacttcccc tatcccaacg tggagaagac tcagaaagac aagaccagtg tgatcaagaa 420
cacagactcc cctgagttca aggagcagtt caaactctgc atcaaccgca gccaccgtgg 480
cttccgaagg ccatccaga ccaagggcat caagtctgaa gtggttcaca agggggggct 540
gttcaagact gaccgggtgc tggggacagc ccagctgaag ctggatgcac tggagatagc 600
atgtgaggtc cgggagatcc ttgaggtcct ggatggtcgc cggccacag gggggcgact 660
ggaggtaatg gtccggatcc gggagccact gacagcccag cagttggaga cgacgacaga 720
gaggtggctg gtcatcgacc ctgtgccggc agctgtgccc acacaggttg ctgggccccaa 780
aggggaaggcc cctcctgtgc ctgcccctgc aagggagtcg gggaaacagat cagccccggcc 840
cctgcatagc ctcatgtgtc tggcggttga ccaagagcgt ctggagcgga agatcctggc 900
cctcaggcag gcgcgccggc cgggtgcccc agaagtggcc cagcagtagc aggacatcat 960
gcaacgcagc cagtggcaga gggcacagct ggagcagggg ggtgtgggca tccgacggga 1020
atacgcagcc cagctggagc ggcagctgca gttctacacg gaggtgccc ggcgcctggg 1080
caacgatggc agcagggatg ctgcaaagga ggcgctctat aggcggaatc tggtagagag 1140
tgagctgcag cggctccgca ggtgaggagc ccatggggcg ggcagcccc agaaagcggg 1200
cagcaggccc cgataccggg aagagccgac acagccacga accagacaag cagacaatca 1260
gcggacaatc ggttctggac tcacccctca tccgggcccc cagccccgcc agagcctccg 1320
tggctgcggg tgttgggaac catgcctgcc agccagtatg tgccctcac ccaggcctgg 1380
ctgggcccctg gagagtcctg tttgcacagc ccaggggtgt ccggcctctg gcccggccc 1440
gagcagggag ggtggctggg gccaaagccc gagggcccct gcaagcactt tacttcctgt 1500
tcctccccag ccttaacccc aaagccctcc tgcaccccaa agaagccact gaggtgggcc 1560
gagccacact gtctccccag gggcgctcgac ctggcccagc tgggtcccca gggccagcac 1620
atggaataaa atagccaggg ccacactcaa aaaaaaaaa 1658

```

&lt;210&gt; 12

&lt;211&gt; 1707

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte Clone No: 2799056

&lt;400&gt; 12

```

ggtgtgcagg atataaggtt ggacttccag acccactgcc cgggagagga gaggagcggg 60
ccgaggactc cagcgtgccc aggtctggca tcctgcactt gctgccctct gacacctggg 120
aagatggccg gcccgaggac cttcaccctt ctctgtggtt tgctggcagc caccttgatc 180
caagccaccc tcagtcccac tgcagttctc atcctcggcc caaaagtcac caaagaaaag 240
ctgacacagg agctgaagga ccacaacgcc accagcatcc tgcagcagct gccgctgctc 300
agtgccatgc gggaaaagcc agccggaggc atccctgtgc tgggcagcct ggtgaacacc 360
gtcctgaagc acatcatctg gctgaaggtc atcacagcta acatcctcca gctgcagggtg 420
aagccctcgg ccaatgacca ggagctgcta gtcaagatcc ccctggacat ggtggctgga 480
ttcaacacgc ccctggtcaa gaccatcgtg gagttccaca tgacgactga ggcccaagcc 540
accatccgca tggacaccag tgcaagtggc cccaccgcc tggtcctcag tgactgtgcc 600
accagccatg ggagcctgcg catccaactg ctgcataagc tctccttcct ggtgaacgcc 660
ttagctaagc aggtcatgaa cctcctagtg ccacccctgc ccaatctagt gaaaaaccag 720
ctgtgtcccg tgatcgaggc ttccttcaat ggcatgtatg cagacctcct gcagctgggtg 780
aaggtgcccc tttccctcag cattgaccgt ctggagtttg accttctgta tcctgccatc 840
aaggtgaca ccattcagct ctacctgggg gccaaagtgt tggactcaca gggaaagggtg 900
accaagtggg tcaataactc tgcagcttcc ctgacaatgc ccaccctgga caacatcccg 960
ttcagcctca tcgtgagtca ggacgtgggtg aaagctgcag tggctgctgt gctctctcca 1020
gaagaattca tggctcctgtt ggactctgtg cttcctgaga gtgcccatcg gctgaagtca 1080
agcatcgggc tgatcaatga aaaggctgca gataagctgg gatctacca gatcgtgaag 1140
atcctaactc aggacactcc cgagtttttt atagaccaag gccatgccaa ggtggcccaa 1200
ctgatcgtgc tggaaagtgt tccctccagt gaagccctcc gccctttgtt caccctgggc 1260
atcgaagcca gctcggaagc tcagttttac accaaagggtg accaacttat actcaacttg 1320
aataacatca gctctgatcg gatccagctg atgaactctg ggattggctg gttccaacct 1380
gatgttctga aaaacatcat cactgagatc atccactcca tcctgctgcc gaaccagaat 1440
ggcaaattaa gatctggggg cccagtgtca ttggtgaagg ccttgggatt cgaggcagct 1500
gagtcctcac tgaccaagga tgcccttggt cttactccag cctccttggt gaaaccagc 1560
tctcctgtct cccagtgaag acttggatgg cagccatcag ggaaggctgg gtcccagttg 1620
ggagtatggg tgtgagctct atagaccatc cctctctgca atcaataaac acttgctgt 1680
gaaaaaaaa aaaaaataaa aaaaaaa 1707

```